

Repeating head-on collisions in an optical trap and the evaluation of spin-dependent interactions among neutral particles

Z B Li, Z F Chen, Y Z He, C G Bao[‡]

State Key Laboratory of Optoelectronic Materials and Technologies, and Department of Physics, Sun Yat-Sen University, Guangzhou, 510275, P.R. China

E-mail: stsbcg@mail.sysu.edu.cn

Abstract. A dynamic process of repeating collisions of a pair of trapped neutral particles with weak spin-dependent interaction is designed and studied. Related theoretical derivation and numerical calculation have been performed to study the inherent coordinate-spin and momentum-spin correlation. Due to the repeating collisions the effect of the weak interaction can be accumulated and enlarged, and therefore can be eventually detected. Numerical results suggest that the Cr-Cr interaction, which has not yet been completely clear, could be thereby determined. The design can be in general used to determine various interactions among neutral atoms and molecules, in particular for the determination of very weak forces.

PACS numbers: 34.90.+q, 34.50.Cx, 03.75.Mn

[‡] The corresponding author

1. Introduction

It is well known that the study of scattering is an important way to understand the interactions among particles. Experimentally, the incident particles with certain momentum would hit a target to initiate the scattering. The accuracy of experimental data depends on a precise control of bombarding energy. For charged incident particles, the initial momenta are usually imposed by electromagnetic forces which can be precisely tuned. For neutral incident particles, the initial momenta are in general difficult to control precisely. Furthermore, the interactions among neutral atoms or molecules are in general weak (say, the van der Waals force). Therefore, the precise determination of these interactions is difficult. Consequently, these interactions are usually determined in an indirect way (say, comparing related experimental data of spectroscopy and/or thermodynamics of a many-body system with theoretical results based on a model with a given set of parameters). In this way the associated theoretical calculations are usually complicated, and uncertainty might exist.

On the other hand, the trapping of neutral atoms has been realized via optical traps since 1998 [1, 2]. This technique is in progress and fewer atoms can be trapped recently [3]. It might open a new way for studying the interactions. In this paper, an idea is proposed and related theoretical calculation is performed to show how the scattering with precisely controllable initial status can be realized in a trap. It turns out that, as we shall see, the collisions would occur repeatedly. Thereby the effect of each individual collision can be accumulated and therefore enlarged. This might lead to a better understanding of very weak interactions among neutral particles.

Since the pioneer experiment by Greismaier, et. al. [4], the Bose-Einstein condensations of atoms with a larger spin (say, ^{52}Cr) become a hot topic. These condensates are a new kind of matter aggregation having the magnetic dipole-dipole interaction V_{dd} more than twenty times stronger than that of the alkalis family.

$$V_{dd} = \frac{C_d}{r^3} \left[\mathbf{F}_1 \cdot \mathbf{F}_2 - 3 \frac{(\mathbf{F}_1 \cdot \mathbf{r})(\mathbf{F}_2 \cdot \mathbf{r})}{r^2} \right], \quad (1)$$

where the strength $C_d = \mu_0 \mu_B^2 g_F^2 / (4\pi)$ with μ_0 being the magnetic permeability of vacuum, μ_B the Bohr magneton, and g_F the Landé g factor, $\mathbf{F}_i$ the operator of the spin of the i -th atom, and $\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$. Consequently, the spatial and spin degrees of freedom are coupled so that the conversion of spin angular momentum into orbital angular momentum can be realized. Thereby new physical phenomena (say, rotonlike behavior) might appear [5, 6]. In addition to the long range V_{dd} , the short range interaction is also spin-dependent and, for low-energy systems, can be in general written as $V_\delta = \delta(\mathbf{r}_1 - \mathbf{r}_2) \sum_S g_S \mathfrak{P}^S$, where S is the total spin, g_S is the strength related to the s -wave scattering length of the S -spin channel, and $\mathfrak{P}^S$ the projector of the S -channel. g_S is nonzero only if S is even. g_2 , g_4 , and g_6 are known while g_0 has not yet [7, 8, 9, 10]. However, many features of the condensate depend strongly on g_0 (say, the phase-diagrams [10, 11, 12, 13] and the spin-evolutions [14]). Therefore, the determination of g_0 is important for a thorough and clear description of this condensate.

As an application of our idea of repeating collisions in a trap, the ^{52}Cr atoms have been chosen as an example to see to what extent the interaction can be thereby clarified.

In the beginning, two narrow and deep potentials $\frac{1}{2}m\omega_a^2|\mathbf{r} \pm \mathbf{a}|^2$ are preset at $\pm \mathbf{a}$ (say, two magnetic traps), where $\mathbf{a}$ is lying along the positive Z -axis. Each potential contains a Cr atom in the ground state of the parabolic confinement. Both atoms are polarized but in reverse directions. The upper (lower) atom has spin-component $\mu = 3$ (-3). Thus the magnetization M_S of the system is zero, and the two atoms are localized. Instantly, the two preset potentials are replaced by a broader potential $\frac{1}{2}m\omega^2 r^2$ (say, an optical trap) centering at the origin. Then, the previously localized atoms begin to evolve. Since $M_S = 0$, the $S = 0$ component must be included and will be affected by g_0 . Therefore, by observing the evolution, the knowledge on g_0 might be extracted.

2. Hamiltonian and initial state

Introduce $\mathbf{R} = (\mathbf{r}_1 + \mathbf{r}_2)/2$ and $\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1$ for the c.m. and relative motions. Introduce $\hbar\omega$ and $\sqrt{\hbar/m\omega}$ as the units of energy and length, respectively. The symmetrized and normalized initial state

$$\Psi_I = \frac{1 + P_{12}}{\sqrt{2}} \left[\left(\frac{2\alpha}{\pi} \right)^{3/4} e^{-\alpha R^2} \cdot \left[\left(\frac{\alpha}{2\pi} \right)^{3/4} e^{-\alpha(r^2/4 + a^2 + ra \cos \theta)} \right] \chi_3(1) \chi_{-3}(2) \right], \quad (2)$$

where P_{12} denotes an interchange of 1 and 2, $\alpha = \omega_a/\omega$, θ is the angle between $\mathbf{r}$ and the Z -axis, $a = |\mathbf{a}|$, and χ_μ is a spin-state of an atom with component μ . Then the evolution is governed by the Hamiltonian

$$H_{evol} = H_R + H_r + V_{12}, \quad (3)$$

where $H_R \equiv -\frac{1}{4}\nabla_R^2 + R^2$ and $H_r \equiv -\nabla_r^2 + \frac{1}{4}r^2$, and $V_{12} = V_\delta + V_{dd}$. The eigenstates of H_R and H_r , denoted as $\bar{\phi}_{NL}(R)Y_{LM}(\hat{R})$ and $\phi_{nl}(r)Y_{lm_l}(\hat{r})$, are just the harmonic oscillator states. The eigenstates of $H_r + V_{12}$ can be expanded in terms of basis functions as

$$\psi_i^J = \sum_{\gamma} C_{i\gamma}^J \phi_{nl}(r) (lS)_J, \quad (4)$$

where γ represents the set n, l , and S , $(lS)_J$ denotes the coupling of l and S into the total angular momentum J , i is just an index of the J -series. Due to V_{dd} , l and S are not conserved, but J is. Due to the boson statistics, $l + S$ must be even.

It turns out that V_{12} is rather weak in our case. Consequently, each eigenstate of $H_r + V_{12}$ is close to an eigenstate of H_r . This fact leads to a great reduction of necessary basis functions in the expansion. When a set of basis functions has been chosen, the associated matrix elements of $H_r + V_{12}$ can be derived as shown in the appendix. Carrying out the diagonalization, the coefficients $C_{i\gamma}^J$ and the corresponding eigenenergy E_i^J can be obtained. In terms of $\bar{\phi}_{NL}$ and ψ_i^J , the initial state can be rewritten as

$$\Psi_I = \sum_N B_N \bar{\phi}_{N,0}(R) Y_{00}(\hat{R}) \sum_{J,i} b_i^J \psi_i^J. \quad (5)$$

Making use of Eq. (4), and equating (5) and (2), it is straight forward to obtain the coefficients B_N and b_i^J .

3. Time-dependent density and the repeating collisions

The time-dependent solution for the evolution is

$$\begin{aligned}\Psi(t) &= e^{-iH_{evol}\tau}\Psi_I \\ &= \frac{1}{\sqrt{4\pi}} \sum_N B_N e^{-i(2N+\frac{3}{2})\tau} \bar{\phi}_{N,0}(R) \\ &\quad \cdot \sum_{J,i} b_i^J e^{-iE_i^J\tau} \psi_i^J,\end{aligned}\tag{6}$$

where $\tau = \omega t$.

From Eq. (6), all the information on the evolution can be extracted. The main feature of the evolution is the occurrence of repeated collisions, as we shall see, and the effect of interaction on each collision can be accumulated and therefore enlarged. Thereby the strength g_0 can be evaluated.

We firstly extract the time-dependent density from $\Psi(t)$ as

$$\begin{aligned}\rho(r, \theta, t) &\equiv 2\pi r^2 \int d\mathbf{R} \Psi^*(t) \Psi(t) \\ &= 2\pi r^2 \sum_{Ji\gamma, J'i'\gamma'} \delta_{S',S} \cos[(E_{i'}^{J'} - E_i^J)\tau] \\ &\quad \cdot b_{i'}^{J'} C_{i'\gamma'}^{J'} b_i^J C_{i\gamma}^J \sum_{M_S} C_{l',-M_S}^{J',0} C_{l,-M_S}^{J,0} \\ &\quad \cdot \phi_{n'l'}(r) \phi_{nl}(r) |Y_{l',-M_S}| |Y_{l,-M_S}|,\end{aligned}\tag{7}$$

where the Clebsch-Gordan coefficients have been introduced. It satisfies

$$1 = \int dr \sin \theta d\theta \rho(r, \theta, t).\tag{8}$$

When the parameters are given as $\omega = 2000 \times 2\pi$, $\alpha = 1.5$, $a = 2$, and $g_0 = -g_6/2$, the variation of ρ in the earliest stage of evolution is shown in Fig. 1. Since ρ does not depend on the azimuthal angle and $\rho(r, \pi - \theta, t) = \rho(r, \theta, t)$, it is sufficient to be plotted only on a quarter of z - x plane, where z and x are the components of $\mathbf{r}/2$. Since the c.m. is always distributed close to the origin, we have roughly $\mathbf{r}_2 \simeq \mathbf{r}/2$ and $\mathbf{r}_1 \simeq -\mathbf{r}/2$. Thus the distribution of an atom can be understood from Fig. 1. When the evolution begins, the two atoms located at opposite ends of the broad potential collide straightly with each other (see 1a and 1b). When $t \approx \pi/2\omega$ (1c), the two atoms keep close to each other with a distance ~ 1.6 , and they are both distributed around the center. Afterward, the atoms begin to separate (1d is very similar to 1b). When $t = \pi/\omega$, the profile (not yet shown) is very similar to 1a. Thus the first round of head-on collision has been completed, and the second round will begin successively. If we remove V_{12} from H_{evol} , the factor $(E_{i'}^{J'} - E_i^J)$ in Eq. (7) would become an integral multiple of $\hbar\omega$, thereby the above process would be exactly periodic with the period $2\pi/\omega$. In fact, the collision as




Figure 1. (Color online) $\rho(r, \theta, t)$ plotted in the z - x plane, where z and x are the components of $\mathbf{r}/2$. t is given at 4 values in the interval (0 to $0.6\pi/\omega = 0.00015$ sec). In the panel, the area with the largest $\rho(r, \theta, t)$ is marked by a $\times$. The parameters are given as $\omega = 2000 \times 2\pi$, $\alpha = 1.5$, $|\mathbf{a}| = 2\sqrt{\hbar/m\omega} = 311$ nm, and $g_0 = -g_6/2$.

shown in Fig. 1 is essentially determined by H_r . In the early stage, V_{12} causes only a very small perturbation. However, as we shall see, the effect of each collision can be accumulated when the time goes on.

The number of collisions that the two atoms have experienced within t is $\sim t\omega/\pi$. When $t = 0.07$ sec and $\omega = 2000 \times 2\pi$, this number is ≈ 280 . When t is close to 0.07 sec, ρ is shown in Fig. 2. At the first glance, 2a is similar to 1a. However, the peak in 2a is considerably lower than that of 1a implying that the atoms are not well localized as in the beginning. In fact, ρ has spread widely in 2a and contains a smooth peak at $x = 1.6$ and $z = 1.4$ (this smooth peak is too low to be seen in the figure). The two atoms are closer to each other in 2b than in 1b. The density varies more vigorously along θ in 2c than in 1c. 2b and 2d do not have the approximate similarity as shown previously in 1b and 1d. From 2a and 2f, we know that a round of collision has been completed in the interval from 0.07 to $0.07 + \pi/\omega$. The evolution in this round is explicitly different from that in the first round due to the accumulated effect of V_{12} .

The accuracy of the above numerical results depends on the number of basis functions, which is determined by the scopes of n (from 0 to $n_{\max}$), l (from 0 to $l_{\max}$), and N (from 0 to $N_{\max}$). When $n_{\max} = N_{\max} = 12$ and $l_{\max} = 14$, the associated results are found to be nearly the same as those by using $n_{\max} = N_{\max} = 10$ and $l_{\max} = 12$. Thus we believe that the former choice is sufficient.




Figure 2. (Color online) The same as Fig. 1 but t is given in the interval $(0.07 \text{ to } 0.07 + \pi/\omega)$.

4. Time-dependent probability of the spin-component and the evaluation

Due to the spin-dependent interaction, spin-flips will occur during the evolution. From Eq. (6) the time-dependent probability of the spin-component of an atom in μ is

$$\begin{aligned}
 P_{\mu}(t) = & \sum_{Ji\gamma, J'i'\gamma'} \delta_{n'n} \delta_{l'l} \cos[(E_{i'}^{J'} - E_i^J)\tau] \\
 & \cdot b_{i'}^{J'} C_{i'\gamma'}^{J'} b_i^J C_{i\gamma}^J \sum_{\lambda} (2\lambda + 1) \sqrt{(2S' + 1)(2S + 1)} \\
 & \cdot W(l3J'3; \lambda S') W(l3J3; \lambda S) \\
 & \cdot C_{\lambda, -\mu}^{J', 0} C_{\lambda, -\mu; 3\mu}^{J, 0},
 \end{aligned} \tag{9}$$

where both the Clebsch-Gordan and Wigner coefficients [15] have been introduced. To show the convergency of numerical calculation, the dependence of $P_{\mu}(t)$ on the number of basis functions is given in Tab. 1.

It is clear from the table that, when $n_{\max} = N_{\max} = 12$ and $l_{\max} = 14$ as we have adopted, qualitatively accurate results can be obtained.

Due to the symmetry inherent in Ψ_I and in H_{evol} , $P_{-\mu}(t) = P_{\mu}(t)$. The variation of $P_3(t)$ against t is shown in Fig. 3, where g_0 is given at three values. It is recalled that the first round of collision ends at $t = \pi/\omega = 0.00025$ sec. In this very short time, the change

Table 1. $P_3(t)$ at three values of t (in sec). The parameters are given as $\omega = 1000 \times 2\pi$, $\alpha = 3$, $|\mathbf{a}| = 2$, and $g_0 = -g_6/2$. The number of basis functions depends on $n_{\max}$ and $l_{\max}$ ($N_{\max} = n_{\max}$ is assumed) listed in the first row.

$n_{\max}$ and $l_{\max}$	10 and 12	12 and 14	14 and 16
$P_3(0.02)$	0.4908	0.4914	0.4914
$P_3(0.06)$	0.4728	0.4733	0.4734
$P_3(0.10)$	0.4751	0.4758	0.4760




Figure 3. (Color online) The evolution of $P_3(t)$. The solid, dash, and dash-dot-dot curves have $g_0 = -g_6/2$, 0, and $g_6/2$, respectively. In 3a, $\omega = 1000 \times 2\pi$, $\alpha = 3$, $|\mathbf{a}| = 2$, while in 3b, $\omega = 2000 \times 2\pi$, $\alpha = 1.5$, $|\mathbf{a}| = 2$.

of $P_3(t)$ is negligible. However, after hundreds of repeating collisions, the dependence of $P_3(t)$ on g_0 can be clearly detected as shown in the figure. Therefore, $P_\mu(t)$ can be used to evaluate g_0 . In particular, the dependence will become more explicit if ω is larger (comparing 3b with 3a). The evolutions of $P_2(t)$ and $P_0(t)$ are shown in Fig. 4a and 4b, respectively, where g_0 is given at seven values. One can see that there is a small peak in 4a appearing in the early stage of evolution if g_0 is negative. Its height depends on how negative g_0 is, and it would disappear if g_0 is positive. Therefore, the existence or not of this peak can be used to judge the sign of g_0 . Together with the height and location of the second peak in Fig. 3 (or the first peak of 4b), the strength g_0 can be known.




Figure 4. (Color online) The evolution of $P_2(t)$ (a) and $P_0(t)$ (b). The curves from "1" to "7" have g_0 from $-3g_6/4$ to $3g_6/4$, respectively, with an increase $g_6/4$ in each step. Each curve has shifted up by 0.005 more than its adjacent lower neighbor to guide the eyes. The other parameters are the same as in Fig. 1.

Since V_{dd} can alter l , the spatial structure can thereby be altered. On the other hand, V_{dd} can also alter S ; therefore spin-evolutions are also affected. We found that, if V_{dd} is reduced (strengthened), the spin-flips would occur less (more) probable. E.g., when V_{dd} is changed to βV_{dd} , the values of the first minimum of the solid curve of Fig. 3b would be 0.462, 0.448, and 0.298, respectively, if $\beta = 0, 1$, and 10. It implies that a strong dipole force will cause strong spin-flips. This is a notable point. On the other hand, recent experimental progress suggests that condensation of molecules with large permanent dipole moments, such as OH [16, 17], RbCs [18], KRb [19], and NH [20], may be achieved. These systems would have very strong dipole interaction, 10^2 or more times stronger than in chromium. Therefore, distinguished phenomena of spins caused by the very strong V_{dd} are expected.

In order to understand better the phenomenon of spin-flips, the momentum-spin correlation is studied in the following. Let the coordinates $\mathbf{r}_1$ and $\mathbf{r}_2$ be introduced via the Talmi-Moshinsky (T-M) coefficients as

$$\begin{aligned} & \bar{\phi}_{N,0}(R)\phi_{nl}(r)Y_{00}(\hat{R})Y_{lm}(\hat{r}) \\ &= \sum_{n_1 l_1 n_2 l_2} a_{n_1 l_1 n_2 l_2}^{N0nl,l} \hat{\phi}_{n_1 l_1}(r_1) \hat{\phi}_{n_2 l_2}(r_2) [Y_{l_1}(\hat{r}_1) Y_{l_2}(\hat{r}_2)]_{lm}, \end{aligned} \quad (10)$$

where l_1 and l_2 are coupled to l and m , and $\hat{\phi}_{nl}(r_i)Y_{lm_l}(\hat{r}_i)$ are the eigenstates of

$H_{r_i} \equiv -\frac{1}{2}\nabla_{r_i}^2 + \frac{1}{2}r_i^2$. The analytical expression of T-M coefficients has been given in the refs. [21, 22, 23]. Inserting Eq. (10) into (6), we have

$$\begin{aligned} \Psi(\tau) = & \sum_{n_1 l_1, n_2 l_2, m_1 m_2, \mu\nu} \delta_{m_1+m_2, -\mu-\nu} Y_{n_1 l_1, n_2 l_2, m_1 m_2}^{\mu\nu}(\tau) \\ & \cdot \hat{\phi}_{n_1 l_1}(r_1) \hat{\phi}_{n_2 l_2}(r_2) Y_{l_1 m_1}(\hat{r}_1) Y_{l_2 m_2}(\hat{r}_2) \\ & \cdot \chi_\mu(1) \chi_\nu(2), \end{aligned} \quad (11)$$

where $Y_{n_1 l_1, n_2 l_2, m_1 m_2}^{\mu\nu}(\tau)$ can be known by comparing Eq. (11) with (6) and (10).

Insert Eq. (11) into the normality $\langle \Psi(\tau) | \Psi(\tau) \rangle = 1$, it is straight forward to obtain the time-dependent probability of finding a particle in χ_ν state at $\mathbf{r}_2$ and τ as follows

$$\begin{aligned} P_\nu(\mathbf{r}_2, \tau) = & \sum_{n'_2 l'_2, n_2 l_2, m_2} Z_{n'_2 l'_2, n_2 l_2, m_2}^\nu(\tau) \hat{\phi}_{n'_2 l'_2}(r_2) \hat{\phi}_{n_2 l_2}(r_2) \\ & \cdot Y_{l'_2 m_2}^*(\hat{r}_2) Y_{l_2 m_2}(\hat{r}_2), \end{aligned} \quad (12)$$

where

$$\begin{aligned} Z_{n'_2 l'_2, n_2 l_2, m_2}^\nu(\tau) = & \sum_{n_1 l_1 m_1 \mu} \delta_{m_1+m_2, -\mu-\nu} [Y_{n_1 l_1, n'_2 l'_2, m_1 m_2}^{\mu\nu}(\tau)]^* \\ & \cdot Y_{n_1 l_1, n_2 l_2, m_1 m_2}^{\mu\nu}(\tau). \end{aligned} \quad (13)$$

This probability fulfills $\sum_\nu \int d\mathbf{r}_2 P_\nu(\mathbf{r}_2, \tau) = 1$.

For observing the momentum-spin correlation, we define

$$\xi_{nl}(p) = \sqrt{\frac{2}{\pi}} (-i)^l \int r^2 dr j_l(pr) \hat{\phi}_{nl}(r), \quad (14)$$

where the spherical Bessel function has been introduced. Then from Eq. (12), we obtain

$$\begin{aligned} Q_\nu(\mathbf{p}, \tau) = & \sum_{n'_2 l'_2, n_2 l_2, m_2} Z_{n'_2 l'_2, n_2 l_2, m_2}^\nu(\tau) \xi_{n'_2 l'_2}^*(p) \xi_{n_2 l_2}(p) \\ & \cdot Y_{l'_2 m_2}^*(\hat{p}) Y_{l_2 m_2}(\hat{p}), \end{aligned} \quad (15)$$

which is the probability of finding a particle in ν at $\mathbf{p}$ and τ . It fulfills $\sum_\nu \int d\mathbf{p} Q_\nu(\mathbf{p}, \tau) = 1$.

In order to observe Q_ν at a given time τ_c , the broad trap has to be suddenly cancelled at τ_c . Then, the particles begin to go out and the successive evolution is an expansion and is governed by the Hamiltonian $H_{free} + V_{12}$, where $H_{free} = -\frac{1}{2}(\nabla_1^2 + \nabla_2^2)$. During the expansion, collisions happen scarcely. We have already seen that the effect of V_{12} within a few collisions is negligible. Therefore, it is safe to neglect V_{12} during the expansion, thus the wave function after τ_c is $\Psi(\tau > \tau_c) \approx e^{-iH_{free}(\tau-\tau_c)} \Psi(\tau_c) \equiv \Psi_{free}(\tau)$.

It is straight forward to see that the weight of $\Psi_{free}(\tau)$ in $|\mathbf{p}_1, \mathbf{p}_2\rangle$ is the same as that of $\Psi(\tau_c)$. Therefore, after the cancellation, the probability Q_ν depends only on τ_c but not on τ . Thus, the time of observation τ is not relevant, but the choice of τ_c is essential.

An example of $Q_3(\mathbf{p}, \tau_c)$ in the earliest stage of evolution is given in Fig. 5a. Curve "1" describes the momentum distribution of the initial h.o. ground state over the Z -component of $\mathbf{p}$. When the evolution begins, the distribution shifts to the left as shown




Figure 5. (Color online) The distribution of $Q_3(\mathbf{p}, \tau_c)$ over the Z -component of $\mathbf{p}$. τ_c is the time to cancel the trap, and is given at five values as $\tau_c = (i - 1)\pi/4 + \tau_X$, "i" is marked by the associated curve, and $\tau_X = 0$ (a) and 200π (b). The unit of momentum in this paper is $\sqrt{m\hbar\omega}$. The parameters are the same as in Fig. 1.

by "2". It implies that the component with $\nu = 3$ is moving from the initial end toward the center. When $\tau = \pi/2$, a sharp peak appears at $p_z = -2$ as shown by "3". Meanwhile, as we know from Fig. 1c, the particle is close to the center, and it has a maximal momentum pointing to the opposite end. Afterward, the magnitude of momentum begins to decrease. When $\tau = \pi$, the average momentum is again zero as shown by "5" which overlaps "1". Then the process proceeds in reverse direction, and repeatedly. When τ is larger, the periodicity will be gradually spoiled by the interaction. This is shown in Fig. 5b, where "5" overlaps "1" no more. Instead, the peak of "5" shifts a little to the left. It implies that the peak has not yet completely arrived at the opposite end, its arrival is delayed.

The momentum-spin correlation can be revealed by $Q_\nu(\mathbf{p}, \tau_c)$. However, we are more interested in a quantity which is easier to be observed. Therefore, we define $Q_\nu^I(\theta_p, \tau_c) \equiv \int p^2 dp Q_\nu(\mathbf{p}, \tau_c)$, where θ_p is the angle between $\mathbf{p}$ and the $+Z$ -axis (the azimuthal angle of $\mathbf{p}$ is irrelevant). This is the probability of finding a particle in ν and emitting at a specified direction θ_p at τ_c . It fulfills $2\pi \int \sin \theta_p d\theta_p Q_\nu^I(\theta_p, \tau_c) = 1$. This quantity is much easier to be experimentally measured because only the orientation of spin is concerned, which can be measured via a Stern-Gerlach device setting towards the outgoing angle θ_p .

Q_ν^I is sensitive to τ_c (therefore both the time of creation and the time of cancellation of the broad potential should be precise). For examples, when $\tau_c = 69.5\pi$ (meanwhile the average positions of both particles are close to the center), Q_3^I against θ_p with g_0 given at three presumed values are plotted in Fig. 6a, where the momentum is distributed around $\theta_p = 0$. Thus the particle is rushing back to the initial end disregarding how g_0 is (meanwhile the $\nu = -3$ particle is rushing to the opposite end). If $\tau_c = 70.5\pi$, the peak of 6a would appear at the right (not yet plotted). When $\tau_c = 70\pi$ (meanwhile the particle is close to the initial end), Q_3^I depends slightly on θ_p as shown in 6b, the dependence is not very sensitive to g_0 .

On the other hand, due to the spin-flips, the $|\nu| \neq 3$ components are created and the creation is sensitive to the interaction. For the case of the $\nu = 2$ component plotted in 6c, the emission with a smaller θ_p would be more probable if g_0 is negative, but less probable if g_0 is positive. The probability $Q_2^I(0, 69.5\pi)$ would decrease by 7 times if g_0 is changed from $-g_6/4$ to $g_6/4$. Thus, information on the interaction can be extracted.

When τ_c increases, Q_2^I would also increase gradually. This is shown in 6c, 6e, and 6f, where $\tau_c = 2k\pi - \pi/2$ (note that the ordinates of these figures have different scales). An example with $\tau_c = 2k\pi$ is shown in 6d. Comparing 6d with 6c, one can see how Q_2^I is changed after the interval $\pi/2$.

The features of other $|\nu| \neq 3$ components are more or less similar to those of $\nu = 2$. Information on the interaction can also be extracted from them. All the curves in Fig. 6 will undergo a left-right reflection if τ_c is changed from $2k\pi - \pi/2$ to $2k\pi + \pi/2$, or if ν is changed to $-\nu$. The reflection is nearly exact.

It is emphasized that the dependence on interaction is hardly found in the early stage. For an example, when $\tau_c = 2\pi$ (meanwhile both particles nearly completely




Figure 6. (Color online) Angular distribution of $Q_\nu^I(\theta_p, \tau_c)$ against θ_p . $\nu = 3$ for (a) and (b), and $\nu = 2$ for the others. τ_c is fixed and marked on the panels. g_0 is given at three values $-g_6/4$, 0 and $g_6/4$, respectively, and the associated curves are in solid, dash, and dash-dot lines. The other parameters are the same as in Fig. 1.

return to their initial positions after two rounds of collisions), the curves of 6b will become a set of horizontal lines and very close to each other, and therefore no valuable information can be extracted. Thus the introduction of the trap, that leads to repeated collisions, is necessary so that the weak effect of interaction can be accumulated.




Figure 7. (Color online) Q_ν^I/Q_3^I with $\theta_p = \pi/2$ and $\tau_c = 69.5\pi$ against g_0/g_6 . ν is marked by the curves. The other parameters are the same as in Fig. 1.

When θ_p and τ_c are both fixed, an example is shown in Fig. 7 where the ratio Q_ν^I/Q_3^I with $\theta_p = \pi/2$ and $\tau_c = 69.5\pi$ is plotted against g_0/g_6 . The curves with ν and $-\nu$ overlap with each other entirely (however, if $\theta_p \neq \pi/2$, they do not). It is clear from the figure that g_0 can be uniquely determined if a few ratios can be measured (say, those with $\nu = 2$ and 0).

5. Conclusions

In summary, an idea is proposed to study the trapped 2-body scattering. The two particles are first localized, then they collide with each other repeatedly in a trap, then they escape. Related theoretical derivation and numerical calculation have been performed to study the coordinate-spin and momentum-spin correlations. The ^{52}Cr atoms have been chosen as an example. Due to the repeating collisions in the trap, the effect of the weak interaction can be accumulated and enlarged, and the strength g_0 can be thereby determined. It is expected that the approach might open a new way

for studying various spin-dependent interactions among atoms (molecules) with nonzero spin, in particular for the studies of very weak forces.

Acknowledgments

The support from the NSFC under the grant 10874249 and from the project of National Basic Research Program of China (2007CB935500) is appreciated.

Appendix A. Matrix elements of V_{12} between the basis functions

$$\begin{aligned} & \langle \phi_{n'l'}(r)(l'S')_{J'} | V_{12} | \phi_{nl}(r)(lS)_J \rangle \\ &= \delta_{J',J} \left[\frac{1}{4\pi} g_S \delta_{l'0} \delta_{l0} \delta_{S'S} \phi_{n'0}(0) \phi_{n0}(0) - 252\sqrt{5} C_d \sqrt{(2S'+1)(2S+1)(2l+1)} \right. \\ & \quad \cdot C_{1,0,1,0}^{2,0} C_{l,0,2,0}^{l',0} W(l2jS'; l'S) \left. \begin{Bmatrix} 1 & 1 & 2 \\ 3 & 3 & S \\ 3 & 3 & S' \end{Bmatrix} \int \frac{dr}{r} \phi_{n'l'}(r) \phi_{nl}(r) \right], \quad (\text{A.1}) \end{aligned}$$

where the Clebsch-Gordan, Wigner and 9- j symbols [15] are introduced. Furthermore, due to the Boson statistics, both $s+l$ and $s'+l'$ should be even.

References

- [1] D. M. Stamper-Kurn, M. R. Andrews, A. P. Chikkatur, S. Inouye, H.-J. Miesner, J. Stenger, and W. Ketterle, Phys. Rev. Lett. **80**, 2027 (1998).
- [2] J. Stenger, S. Inouye, D. M. Stamper-Kurn, H.-J. Miesner, A. P. Chikkatur, and W. Ketterle, Nature (London) **396**, 345 (1998).
- [3] M. Albiez, R. Gati, J. Fölling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler, Phys. Rev. Lett. **95**, 010402 (2005).
- [4] A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, Phys. Rev. Lett. **94**, 160401 (2005).
- [5] D. H. J. O'Dell, S. Giovanazzi, and G. Kurizki, Phys. Rev. Lett. **90**, 110402 (2003).
- [6] L. Santos, G. V. Shlyapnikov, and M. Lewenstein, Phys. Rev. Lett. **90**, 250403 (2003).
- [7] J. Werner, A. Griesmaier, S. Hensler, J. Stuhler, T. Pfau, A. Simoni, and E. Tiesinga, Phys. Rev. Lett. **94**, 183201 (2005).
- [8] J. Stuhler, A. Griesmaier, T. Koch, M. Fattori, T. Pfau, S. Giovanazzi, P. Pedri, and L. Santos, Phys. Rev. Lett. **95**, 150406 (2005).
- [9] A. Griesmaier, J. Stuhler, T. Koch, M. Fattori, T. Pfau, and S. Giovanazzi, Phys. Rev. Lett. **97**, 250402 (2006).
- [10] R. B. Diener and T. L. Ho, Phys. Rev. Lett. **96**, 190405 (2006).
- [11] H. Mäkelä and K.-A. Suominen, Phys. Rev. A **75**, 033610 (2007).
- [12] S. Uchino, T. Otsuka, and M. Ueda, Phys. Rev. A **78**, 023609 (2008).
- [13] P. Van Isacker and S. Heinze, J. Phys. A **40**, 14811 (2007).
- [14] L. Santos and T. Pfau, Phys. Rev. Lett. **96**, 190404 (2006).
- [15] A. R. Edmonds, *Angular momentum in quantum mechanics* (Princeton, Princeton University Press, 1957).
- [16] S. Y. T. van de Meerakker, P. H. M. Smeets, N. Vanhaecke, R. T. Jongma, and G. Meijer, Phys. Rev. Lett. **94**, 023004 (2005).
- [17] J. R. Bochinski, E. R. Hudson, H. J. Lewandowski, and J. Ye, Phys. Rev. A **70**, 043410 (2004).

- [18] J. M. Sage, S. Sainis, T. Bergeman, and D. DeMille, Phys. Rev. Lett. **94**, 203001 (2005).
- [19] D. Wang, J. Qi, M.F. Stone, O. Nikolayeva, B. Hattaway, S.D. Gensemer, H. Wang, W.T. Zemke, P.L. Gould, E.E. Eyler and W.C. Stwalley, Eur. Phys. J. D **31**, 165 (2004).
- [20] D. Egorov, W.C. Campbell, B. Friedrich, S.E. Maxwell, E. Tsikata, L.D. van Buuren and J.M. Doyle, Eur. Phys. J. D **31**, 307 (2004).
- [21] W. Tobocman, Nucl. Phys. A. **357**, 293 (1981).
- [22] M. Baranger, and K. T. R. Davies, Nucl. Phys. **79**, 403 (1966)
- [23] T. A. Brody and M. Moshinski, Monografias del Instituto de Fisica, Universidad Nacional Autonoma de Mexico (1960).